

ACTION

lidafol KL is a liquid foliar fertilizer specially formulated for use during the ripening stage.

lidafol KL is intended to supplement standard fertility programs and is a readily available source of potassium, phosphorus, boron and molybdenum.

CAUTION

lidafol KL is recommended only for use in citrus (lemon, orange, mandarin and clementine) vegetables (cucurbits, pepper, tomato, aubergine (eggplant), strawberry, beans, artichoke, and potato), tree fruit (apples, cherry, loquat, nectarines, peaches, and plum) and grapes (table and wine).

The use of **lidafol KL** on any other crop or at any other timing not listed on the label may result in serious crop injury. The application of fertilizing materials containing molybdenum (Mo) may result in forage crops containing levels of molybdenum (Mo) which are toxic to ruminant animals.

WARNINGS

USE ONLY IN CASE OF NECESSITY.

DO NOT EXCEED THE RECOMMENDED RATE.

The recommendations and information given in this label have been carefully studied and tested. However, in the utilization of the product, there could be interference in other factors that are not possible to control such as mixture preparation, application, climate, etc. Lida Plant Research LLC. guarantees the composition, formulation and content. The user will be responsible for the damages caused by lack of efficacy, toxicity, residues, etc., by partial and/or total un-observance of the instructions given on the label. Keep out of reach of children. Keep away from food, drinks and cattle food.

Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapfco.org/metals.htm>

plant nutrition

lidafol KL



0-15-20

GUARANTEED ANALYSIS

| | |
|--|-------|
| Available Phosphate (P ₂ O ₅) | 15 % |
| Soluble Potash (K ₂ O) | 20 % |
| Boron (B) | 0.2 % |
| Molybdenum (Mo) | 0.2 % |

Derived from dipotassium phosphate, boric acid and ammonium heptamolybdate.

Manufactured by:



Guaranteed/Distributed by:

Lida Plant Research LLC.
1444 Fulton Street
Fresno, CA 93721
Phone (559) 5003535
Fax (559) 5003536

CROPS

Citrus: Especial for early varieties and fine lemon, as well as for all the varieties of orange, mandarin and Clementine.

Vegetables: Very indicated for forced crops under plastics, especially for water melon, melon, pepper, tomato, aubergine, strawberry, stick bean, artichoke and potato.

Fruit trees: Nectarine, peach, grape, loquat, apple, plum and cherry trees

Wine grapes and table grapes.

FOLIAR DOSE

0.2 - 0.3 % solution

In citrus, fruit trees and vines apply at least 2 treatments and in exceptional cases 3 treatments from the start of the fruit coloring until the final ripening.

In vegetables apply several treatments throughout the cultivation cycle, every 10 - 12 days, starting to treat as from the fruit differentiation.

STORAGE

Avoid low temperatures in winter and temperatures above 82.4°F/28°C.

Avoid direct light and long storage periods.

Density: 12.52 lbs/gal (1.5 kg/L) at 68 °F

Net Volume: 1.32 gals (5 Liters) or 0.26 gals (1 Liter)

Net Weight 16.52 lbs (7.5 kg) or 3.30 lbs (1.5 kg)