



trevian®

1.6 - 5 - 3.2

GUARANTEED ANALYSIS

Total Nitrogen (N)	1.6 %
1.6 % Water Soluble Nitrogen	
Available Phosphate (P2O5)	5 %
Soluble Potash (K2O)	3.2 %

Derived from seafood by-product (hydrolysate from crab shells and shrimps), and monopotassium phosphate

ACTION

Trevian is recommended for use as a supplement to improve crop productivity and promote plant vigor in agricultural crops.

STORAGE

Avoid low temperatures in winter and temperatures above 82°F/28°C. Avoid direct light and long storage periods.

WARNINGS

**USE ONLY IN CASE OF NECESSITY.
DO NOT EXCEED THE RECOMMENDED RATE.**

The recommendations and information given in this label have been carefully studied and tested. However, when applying the product, there could be interferences with other factors that are not possible to control such as mixture preparation, application, climate, etc. Lida Plant Research LLC. guarantees the composition, formulation and content. The user is responsible for any damages caused by lack of efficacy, toxicity, residues, etc. by partial and/or total un-observance of the instructions given on the label. Keep out of reach of children. Keep away from food, drinks and cattle feed.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

Manufactured by:
Guaranteed/Distributed by:
Lida Plant Research LLC.
1444 Fulton Street
Fresno, CA 93721
Phone (559) 5003535
Fax (559) 5003536



CROPS AND DOSE

NURSERIES AND POTTED PLANTS

- Foliar spray: 8 - 16 oz/ac.
- Fertiligation: 16 oz/ac.

CEREALS, ROW CROPS, AND GRASSES: Corn, Cotton, Rice, Wheat, Barley, Bluegrass (For Seed): 1-3 applications per season. Apply every 21 from first true leaves.

- Foliar spray: 6 - 8 oz/ac.

VEGETABLES: Beet, Broccoli, Brussel Sprouts, Cabbage, Carrot, Cauliflower, Celery, Garlic, Green Beans, Kale, Leafy Greens, Lettuce, Melon, Onion, Pepper, Potatoe, Pumpkin, Radish, Spinach, Tomato, Watermelon: 3-4 applications per season. Apply every 10-15 days throughout the growing season.

- Foliar spray: 8 - 16 oz/ac.
- Fertiligation: 16 oz/ac.

BERRIES: Blackberry, Blueberry, Raspberry, Strawberry: 4-5 Application per season. Apply every 10-15 days throughout the growing season.

- Foliar spray: 12 - 16 oz/ac.
- Fertiligation: 16 oz/ac

FRUIT TREES: Persimmon, Grapes (Table and Wine), Stonefruits (Apricot, Cherry, Nectarine, Peach), Nuts (Almond, Pecan, Pistachio, Walnut) 2-3 apply every 15-21 days making the first application during the fruit setting.

- Foliar spray: 16 oz/ac

In general, for terrestrial application check minimum and maximum concentrations: 1.5 to 2 oz each 10 gal of water.

Net Weight: 2 lbs. (0.9 kg.)