

## ACTION

**vaccilan** is a fertilizer designed to supply potassium as a foliar spray on most crops.

## WARNINGS

### DO NOT EXCEED THE RECOMMENDED RATE.

The recommendations and information given in this label have been carefully studied and tested. However, when applying the product, there could be interferences with other factors that are not possible to control such as mixture preparation, application, climate, etc. Lida Plant Research LLC. guarantees the composition, formulation and content. The user is responsible for any damages caused by lack of efficacy, toxicity, residues, etc. by partial and/or total un-observance of the instructions given on the label. Keep out of reach of children. Keep away from food, drinks and cattle feed.

Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapfco.org/metals.html>

## STORAGE

Avoid low temperatures in winter and temperatures above 82°F/28°C. Avoid direct light and storage longer than 2 years.

## CROPS AND RATES

**VEGETABLES:** fruit (pepper, tomato, melons, cucumber, squash, pumpkin, green beans, green peas), onion, garlic, potato, sweet potato, carrot, beets, cabbage, cauliflower, broccoli, brusel sprouts, radish, pak choy, bok choy, pe-tsai, other brassicas, leafy greens (lettuce, celery, coriander (cilantro), parsley, chards, spinach), asparagus, artichoke

**Foliar rates:** 8 - 20 fl oz/ac.

**Directions:** For spray volume less than 30 gpa use lower rates and/or concentrations of 3 to 4 fl oz each 10 gal. For spray volume above 50 gpa use higher rates and/or concentrations of 2 to 3 fl oz each 10 gal. For aircraft applications use low to medium rate. It is recommended to do the first two applications every 7 days and the next two every 21 days.



# vaccilan

## 0-0-3

## GUARANTEED ANALYSIS

Soluble Potash (K<sub>2</sub>O)

3 %

Derived from Potassium formate

Manufactured by:  
LIDA PLANT RESEARCH  
Spain (EUROPEAN UNION)



Guaranteed/Distributed by:  
Lida Plant Research LLC.  
1444 Fulton Street  
Fresno, CA 93721  
Phone (559) 5003535  
Fax (559) 5003536

**Density:** 8.76 lbs./gal (1.05 kg/L) at 68 °F.  
**Net Volume:** 0.25 gals (0.95 L)  
**Net Weight:** 2.2 lbs. (1 Kg)

## CROPS AND RATES

**STRAWBERRY, BLACKBERRY, RASPBERRY, BLUEBERRY:** **Foliar rates:** 16 - 32 fl oz/ac **Directions:** For spray volume less than 100 gpa use low to medium rates and/or concentrations of 1.5 to 3 fl oz each 10 gal. For spray volume above 120 gpa use medium to high rates and/or concentrations of 1 to 1.5 fl oz each 10 gal. For aircraft applications use low to medium rate. It is recommended to do the first two applications every 7 days and the next two every 21 days.

**VINEYARD (WINE AND TABLE):** **Foliar rates:** 16 - 32 fl oz/ac **Directions:** For spray volume less than 80 gpa use low to medium rates and/or concentrations of 1.5 to 3 fl oz each 10 gal. For spray volume above 120 gpa use medium to high rates and/or concentrations of 1 to 2 fl oz each 10 gal. For aircraft applications use low to medium rate. It's recommended to do 2-3 applications, with 21 days between applications.

**FRUIT AND NUT TREES:** orange, lemon, lime, mandarin (and any kind of hybrids), grapefruit, apple, stonefruits (peach, apricot, nectarine, plum, prune, cherry), nuts (almond, hazelnut), avocado, fig. **Foliar rates:** 16 - 32 fl oz/ac **Directions:** For spray volume less than 80 gpa use low to medium rates and/or concentrations of 1.5 to 3 fl oz each 10 gal. For spray volume above 120 gpa use medium to high rates and/or concentrations of 1 to 2 fl oz each 10 gal. For aircraft applications use low to medium rate. It's recommended to do 2-3 applications, with 21 days between applications.

**ROW CROPS:** corn, sorghum, wheat, barley, dry beans, soy beans, cotton, sugarbeets and **FORAGE CROPS:** hay grasses: **Foliar rates:** 4 - 10 fl oz/ac **Directions:** For terrestrial or aircraft applications. For terrestrial applications use between 2 and 4 fl oz each 10 gal/ac of spray volume. It's recommended to do 1-2 applications, with 20-25 days between applications

**NURSERIES & LANDSCAPE:** ornamental, trees, turf, sod **Foliar rates:** 8 - 20 fl oz/ac **Directions:** For spray volume less than 30 gpa use lower rates and/or concentrations of 3 to 4 fl oz each 10 gal. For spray volume above 50 gpa use higher rates and/or concentrations of 2 to 3 fl oz each 10 gal. It's recommended to do 2-3 applications, with 15-21 days between applications.

**HYDROPONIC CROPS:** For foliar rates use the recommended rate on this label for the specific commodity.

**DO NOT EXCEED THE INDICATED APPLICATION RATES. USE ONLY ON CROPS INDICATED ON THE LABEL.**